



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

From (1), $z=.00066$, $s=6416.9M$. The ultimate strength of hard steel is 240000 lb./in.².

$$\therefore s=240000/f=240000/6=40000 \text{ lb./in.}^2$$

$$\therefore 40000=6416.9M. \quad \therefore M=6.2335.$$

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

159. Proposed by E. B. ESCOTT, Ann Arbor, Mich.

Show that if the equation $y^3=2x^2-1$ be possible in integers, $y=24n^2-1$, or $2n^2-1$, and find three solutions.

Solution by the PROPOSER.

The equation may be written $(y+1)(y^2-y+1)=2x^2$.

Since y^2-y+1 is always odd, it is evident that $y+1$ must be even. Since $y^2-y+1=(y+1)(y-2)+3$ it is evident that $y+1$ and y^2-y+1 can have no common factor but 3. Therefore we have the following possibilities for y : $y=2 \times 3m^2-1$, or $y=2n^2-1$.

Since y^3 is represented by the form $2x^2-1$, 2 must be a quadratic residue of y . Therefore $y=8a \pm 1$, and this is possible in the first expression only when $m=2n$. Then either $y=24n^2-1$ or $2n^2-1$.

Substituting these values of y in the original equation, we have

$$192n^4-24n^2+1=r^2 \text{ or } 4n^4-6n^2+3=r^2.$$

The first equation has the solution $n=0$ and 1 which give $y=-1$, $x=0$; $y=23$, $x=78$.

The second equation has the solution $n=1$, which gives $y=1$, $x=1$.

AVERAGE AND PROBABILITY.

198. Proposed by J. EDWARD SANDERS, Weather Bureau, Chicago, Ill.

Find the average length of a hole at random through a given cylinder.

No solution of this problem has been received.

199. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

A circle is inscribed in a given square. Two points are taken at random within the square but without the circle. What is the chance the distance between the points does not exceed the side of the square?

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

This is the same as 196, but as there is a distance less than the side of the square when both points are taken one each in opposite corners, it is